

Exam

A number of different exercises is proposed. You do not need to try to solve them all.

Exercise 1 Let R be a relation on a set X , that is, a subset of X^2 . Let R^0 be the identity relation, that is, x is in relation with y , written xR^0y , if $x = y$. And let R^{n+1} be the composition of R and R^n . The composition of two relations R and S , written $R \circ S$, is the relation such that $x(R \circ S)y$ if there is z such that xRz and zRy . Hence, xR^ny ($n > 0$) if there are $n - 1$ intermediate elements x_1, \dots, x_{n-1} such that $xRx_1, \dots, x_{n-1}Ry$. A relation S is reflexive if, for all x , xSx . A relation S is transitive if, for all x, y, z , xSz whenever xSy and ySz . Prove that:

1. $R^+ = \bigcup_{i>0} R^i$ is transitive.
2. R^+ is the smallest transitive relation containing R .
3. $R^* = \bigcup_{i\geq 0} R^i$ is transitive and reflexive.
4. R^* is the smallest transitive and reflexive relation containing R .
5. The function $f(X) = X \circ X$ is monotone, that is, $f(X) \subseteq f(Y)$ if $X \subseteq Y$.
6. R^* is a fixpoint of f , that is, $f(R^*) = R^*$.

Exercise 2 We consider the set of λ -terms $t = x \mid \lambda xt \mid tt$, where x is a variable taken in an infinite set \mathcal{X} . Let \rightarrow_β be the smallest relation such that:

- $(\lambda xt)u \rightarrow_\beta t\{x \mapsto u\}$, where $t\{x \mapsto u\}$ is the term obtained by replacing every free occurrence of x in t by u (and renaming bound variables if necessary)
- $\lambda xt \rightarrow_\beta \lambda xt'$ if $t \rightarrow_\beta t'$
- $tu \rightarrow_\beta t'u$ if $t \rightarrow_\beta t'$
- $tu \rightarrow_\beta tu'$ if $u \rightarrow_\beta u'$

Prove that:

1. β -reduction introduces no new variables: if $t \rightarrow_\beta t'$, then $\text{FV}(t') \subseteq \text{FV}(t)$.
2. If $v \rightarrow_\beta^* v'$, then $t\{y \mapsto v\} \rightarrow_\beta^* t\{y \mapsto v'\}$.

3. \rightarrow_β is stable by substitution: if $t \rightarrow_\beta t'$, then $t\{y \mapsto v\} \rightarrow_\beta t'\{y \mapsto v\}$.
4. \rightarrow_β is locally confluent, that is, if $t \rightarrow_\beta u$ and $t \rightarrow_\beta v$, then there is w such that $u \rightarrow_\beta^* w$ and $v \rightarrow_\beta^* w$.

This last property means that, locally, the order in which we make β -reductions is not important, since we can always find a common reduct. An important property of β -reduction is that it also holds globally: if $t \rightarrow_\beta^* u$ and $t \rightarrow_\beta^* v$, then there is w such that $u \rightarrow_\beta^* w$ and $v \rightarrow_\beta^* w$. Thus, β -reduction is in some sense deterministic: it can lead to only one result.

Exercise 3 Define a 2-tapes Turing machine $(\Sigma, Q, q_i, q_f, \delta)$ computing the function $n \mapsto 2n$, where Σ is the finite alphabet used by the machine with at least the symbols $\{\square, 0, 1\}$, Q is the finite set of states of the machine, q_i the initial state, q_f the final state, and δ is a partial function from $Q \times \Sigma^2$ to $Q \times \Sigma^2 \times \{-1, 0, +1\}$. To do this, you are free to introduce in Σ symbols different from $\{\square, 0, 1\}$, and in Q states different from $\{q_i, q_f\}$. The machine starts in position 0 in state q_i with, on tape 1, \square followed by n 1's and then 0's, and on tape 2, \square followed by 0's. It must end in position 0 in state q_f with, on tape 2, \square followed by $2n$ 1's and then 0's.

Exercise 4 We consider a set \mathcal{X} of variables, a set \mathcal{F} of function symbols and a set \mathcal{P} of predicate symbols. We assume that every (function or predicate) symbol f is equipped with an arity $\alpha_f \in \mathbb{N}$, that is, in a term of the form $f(t_1, \dots, t_n)$, we always have $n = \alpha_f$. A model is given by a set A and, for each function symbol f of arity n , a function $f_A : A^n \rightarrow A$ and, for each predicate symbol P of arity n , a function $P_A : A^n \rightarrow \text{Bool}$ where $\text{Bool} = \{\text{true}, \text{false}\}$. Then, given a finite valuation $\mu : \mathcal{X} \rightarrow A$, the interpretation of a formula ϕ , written $\llbracket \phi \rrbracket_\mu$, is defined as follows:

- $\llbracket x \rrbracket_\mu = \mu(x)$
- $\llbracket f(t_1, \dots, t_n) \rrbracket_\mu = f_A(\llbracket t_1 \rrbracket_\mu, \dots, \llbracket t_n \rrbracket_\mu)$
- $\llbracket P(t_1, \dots, t_n) \rrbracket_\mu = P_A(\llbracket t_1 \rrbracket_\mu, \dots, \llbracket t_n \rrbracket_\mu)$
- $\llbracket \perp \rrbracket_\mu = \text{false}$
- $\llbracket \neg \phi \rrbracket_\mu = \text{not}(\llbracket \phi \rrbracket_\mu)$
- $\llbracket \phi \vee \psi \rrbracket_\mu = \text{or}(\llbracket \phi \rrbracket_\mu, \llbracket \psi \rrbracket_\mu)$
- $\llbracket \phi \wedge \psi \rrbracket_\mu = \text{and}(\llbracket \phi \rrbracket_\mu, \llbracket \psi \rrbracket_\mu)$
- $\llbracket \phi \Rightarrow \psi \rrbracket_\mu = \text{impl}(\llbracket \phi \rrbracket_\mu, \llbracket \psi \rrbracket_\mu)$
- $\llbracket \forall x \phi \rrbracket_\mu = \text{forall}(\{\llbracket \phi \rrbracket_{\mu \cup \{x, a\}} \mid a \in A\})$ if $x \notin \text{dom}(\mu)$
- $\llbracket \exists x \phi \rrbracket_\mu = \text{exists}(\{\llbracket \phi \rrbracket_{\mu \cup \{x, a\}} \mid a \in A\})$ if $x \notin \text{dom}(\mu)$

where $\text{dom}(\mu)$ is the set of variables on which μ is defined, the boolean functions **not**, **or**, **...** are defined as usual, and **forall**, **exists** : $\mathcal{P}(\text{Bool}) \rightarrow \text{Bool}$ are defined as follows:

- $\text{forall}(S) = \text{true}$ iff $\text{false} \notin S$
- $\text{exists}(S) = \text{true}$ iff $\text{true} \in S$

The universal closure of a formula ϕ with free variables x_1, \dots, x_n is $\bar{\forall}\phi = \forall x_1 \dots \forall x_n \phi$. A formula ϕ is *valid* if, in every model A , $\llbracket \bar{\forall}\phi \rrbracket = \text{true}$.

Let the provability relation \vdash be the smallest relation on pairs (Γ, ϕ) where Γ is a finite set of formulas (the assumptions) and ϕ a formula, such that:

$$\begin{aligned}
(\text{axiom}) \quad & \frac{\phi \in \Gamma}{\Gamma \vdash \phi} \\
(\perp\text{-elim}) \quad & \frac{\Gamma \vdash \perp}{\Gamma \vdash \phi} \\
(\Rightarrow\text{-intro}) \quad & \frac{\Gamma \cup \{\phi\} \vdash \psi}{\Gamma \vdash \phi \Rightarrow \psi} \\
(\Rightarrow\text{-elim}) \quad & \frac{\Gamma \vdash \phi \Rightarrow \psi \quad \Gamma \vdash \phi}{\Gamma \vdash \psi} \\
(\wedge\text{-intro}) \quad & \frac{\Gamma \vdash \phi \quad \Gamma \vdash \psi}{\Gamma \vdash \phi \wedge \psi} \\
(\wedge\text{-elim-left}) \quad & \frac{\Gamma \vdash \phi \wedge \psi}{\Gamma \vdash \phi} \\
(\wedge\text{-elim-right}) \quad & \frac{\Gamma \vdash \phi \wedge \psi}{\Gamma \vdash \psi} \\
(\vee\text{-intro-left}) \quad & \frac{\Gamma \vdash \phi}{\Gamma \vdash \phi \vee \psi} \\
(\vee\text{-intro-right}) \quad & \frac{\Gamma \vdash \psi}{\Gamma \vdash \phi \vee \psi} \\
(\vee\text{-elim}) \quad & \frac{\Gamma \vdash \phi \vee \psi \quad \Gamma \cup \{\phi\} \vdash \chi \quad \Gamma \cup \{\psi\} \vdash \chi}{\Gamma \vdash \chi} \\
(\forall\text{-intro}) \quad & \frac{\Gamma \vdash \phi \quad x \notin \text{FV}(\Gamma)}{\Gamma \vdash \forall x \phi} \\
(\forall\text{-elim}) \quad & \frac{\Gamma \vdash \forall x \phi}{\Gamma \vdash \phi\{x \mapsto t\}}
\end{aligned}$$

$$(\exists\text{-intro}) \frac{\Gamma \vdash \phi\{x \mapsto t\}}{\Gamma \vdash \exists x\phi}$$

$$(\exists\text{-elim}) \frac{\Gamma \vdash \exists x\phi \quad \Gamma \cup \{\phi\} \vdash \chi \quad x \notin \text{FV}(\Gamma) \cup \text{FV}(\chi)}{\Gamma \vdash \chi}$$

$$(\text{EM}) \Gamma \vdash \phi \vee \neg\phi$$

Prove that the provability relation is correct, that is, in every model A , if $\Gamma \vdash \phi$ and $\Gamma = \{\phi_1, \dots, \phi_n\}$, then $\phi_1 \wedge \dots \wedge \phi_n \Rightarrow \phi$ is valid.